

VIDYA BHAWAN BALIKA VIDYA PITH
शक्तिउत्थानआश्रमलखीसरायबिहार

Class 12 Sub-.Maths

Date 10.06..2021

1. A spherical ball of salt is dissolving in water in such a manner that the rate of decrease of the volume at any instant is proportional to the surface. Prove that the radius is decreasing at a constant rate.

Solution:

Given, a spherical ball of salt

Then, the volume of ball $V = \frac{4}{3} \pi r^3$ where r = radius of the ball

Now, according to the question we have

$dV/dt \propto S$, where S = surface area of the ball

$$\frac{d}{dt} \left(\frac{4}{3} \pi r^3 \right) \propto 4\pi r^2 \quad [\because S = 4\pi r^2]$$

$$\frac{4}{3} \pi \cdot 3r^2 \cdot \frac{dr}{dt} \propto 4\pi r^2$$

$$4\pi r^2 \cdot \frac{dr}{dt} = K \cdot 4\pi r^2 \quad (K = \text{Constant of proportionality})$$

$$\frac{dr}{dt} = K \cdot \frac{4\pi r^2}{4\pi r^2}$$

$$\frac{dr}{dt} = K \cdot 1 = K$$

Therefore, the radius of the ball is decreasing at constant rate.

2. If the area of a circle increases at a uniform rate, then prove that perimeter varies inversely as the radius.

Solution:

We know that the area of circle, $A = \pi r^2$, where r = radius of the circle

And, perimeter = $2\pi r$

According to the question, we have

$$\frac{dA}{dt} = K, \text{ where } K = \text{constant}$$

$$\Rightarrow \frac{d}{dt}(\pi r^2) = K \Rightarrow \pi \cdot 2r \cdot \frac{dr}{dt} = K$$

$$\text{So, } \frac{dr}{dt} = \frac{K}{2\pi r}$$

Now, perimeter $c = 2\pi r$

Differentiating w.r.t t , we get

$$\frac{dc}{dt} = \frac{d}{dt}(2\pi r) \Rightarrow \frac{dc}{dt} = 2\pi \cdot \frac{dr}{dt}$$

$$\frac{dc}{dt} = 2\pi \cdot \frac{K}{2\pi r} = \frac{K}{r}$$

[From (1)]

$$\frac{dc}{dt} \propto \frac{1}{r}$$


Therefore, it's seen that the perimeter of the circle varies inversely as the radius of the circle.

3. A kite is moving horizontally at a height of 151.5 meters. If the speed of kite is

10 m/s, how fast is the string being let out; when the kite is 250 m away from the boy who is flying the kite? The height of boy is 1.5 m.

Solution:

Given,



Height of the kite(h) = 151.5 m

Speed of the kite(V) = 10 m/s

Let FD be the height of the kite and AB be the height of the kite and AB be the height of the boy.

Now, let AF = x m

So, BG = AF = x

And, dx/dt = 10 m/s

From the figure, it's seen that

$$GD = DF - GF = DF - AB$$

$$= (151.5 - 1.5) \text{ m} = 150 \text{ m [As AB = GF]}$$

Now, in ΔBDG

$$BG^2 + GD^2 = BD^2 \text{ (By Pythagoras Theorem)}$$

$$x^2 + (150)^2 = (250)^2$$

$$x^2 + 22500 = 62500$$

$$x^2 = 62500 - 22500 = 40000$$

$$x = 200 \text{ m}$$

Let initially the length of the string be y m

So, in ΔBDG

$$BG^2 + GD^2 = BD^2$$

$$x^2 + (150)^2 = y^2$$

Differentiating both sides w.r.t., t, we have

$$2x \cdot \frac{dx}{dt} + 0 = 2y \cdot \frac{dy}{dt} \quad \left[\because \frac{dx}{dt} = 10 \text{ m/s} \right]$$
$$2 \times 200 \times 10 = 2 \times 250 \times \frac{dy}{dt}$$
$$\frac{dy}{dt} = \frac{2 \times 200 \times 10}{2 \times 250} = 8 \text{ m/s}$$

Therefore, the rate of change of the length of the string is 8 m/s.