
## CHEMISTRY STUDY MATERIALS FOR CLASS 12 (NCERT Based Notes of Chapter - 12) GANESH KUMAR DATE:- 15/11/2020

# Aldehyde, Ketones and Carboxylic Acid

## 5. Aldol condensation Reaction:

Aldehydes and ketones having at least one  $\alpha$ -hydrogen atom when treated with dilute alkali, we get  $\beta$ -hydroxy aldehydes (aldol) or  $\beta$ -hydroxy ketones (ketol) respectively. This is known as Aldol reaction. The product formed contains both aldehydic (ketonic) and alcoholic group. So the name aldol or ketol is used.


The aldol or ketol on heated undergo dehydration to give  $\alpha$ ,  $\beta$ -unsaturated aldehyde or ketone. This reaction is called Aldol condensation.



6. Cross aldol condensation: When aldol condensation is carried out between two different aldehydes or ketones, it is called cross aldol condensation. If both of them contain α-hydrogen atoms, we get a mixture of four products.
CH<sub>3</sub>-CHO + CH<sub>2</sub>-CH<sub>2</sub>-CHO \_\_\_\_\_\_ CH<sub>3</sub>-CH=CH-CHO +

 $CH_3-CHO + CH_2-CH_2-CHO \longrightarrow CH_3-CH=CH-CHO +$ Ethanal Propanal But-2-enal


 $CH_3-CH_2-CH=C(CH_3)-CHO + CH_3-CH=C(CH_3)-CHO + CH_3-CH_2-CH=CH-CH$ 2-methylpent-2-enal 2-methylbut-2-enal Pent-2-enal



1, 3-Diphenylprop-2-en-1-one (Benzalacetophenone)

## 7. Cannizzaro Reaction:


Aldehydes having no  $\alpha$ -hydrogen atom (e.g. HCHO, C<sub>6</sub>H<sub>5</sub>-CHO, CCl<sub>3</sub>-CHO etc), when treated with Conc. alkali (NaOH or KOH) undergo self oxidation and reduction (disproportionation) to form one molecule of the alcohol and one molecule of carboxylic acid salt. This reaction is called Cannizzaro reaction.



### 8. Electrophilic Substitution Reactions:

Aldehydic and ketonic groups are meta directing and deactivating. So on electrophilic substitution reactions, they give meta-derivatives.

#### e.g. Nitration:



Benzaldehyde

*m*-Nitrobenzaldehyde

### Tests to distinguish Aldehydes and Ketones

#### 1. Tollens' test:

Tollen's reagent is freshly prepared *ammoniacal Silver nitrate*. On warming Tollens' reagent, aldehydes give a bright silver mirror. During this reaction, the aldehyde is oxidised to corresponding carboxylate ion and silver nitrate is reduced to silver metal.

$$R-CHO + 2[Ag(NH_3)_2]^+ + 3OH^- \rightarrow R-COO^- + 2Ag + 2H_2O + 4NH_3$$

#### 2. Fehling's test:

Fehling reagent is a mixture of two solutions, Fehling solution A and Fehling solution B. Fehling solution A is *aqueous copper sulphate* and Fehling solution B is *alkaline sodium potassium tartarate (Rochelle salt)*. On heating with Fehling's reagent, aldehyde gives a reddish brown precipitate of cuprous oxide (Cu<sub>2</sub>O). Aromatic aldehydes do not give this test.

 $R-CHO + 2Cu^{2+} + 5OH^{-} \longrightarrow R-COO^{-} + Cu_2O + 3H_2O$ 

(Cuprous oxide)

#### 3. Benedict's test:

Benedict's solution is a mixture of *CuSO*<sub>4</sub>, sodium citrate and sodium carbonate. Aliphatic aldehydes give reddish brown precipitate when heated with Benedict's reagent.

 $R-CHO + 2Cu^{2+} + 5OH^{-}$   $R-COO^{-} + Cu_{2}O + 3H_{2}O$ 

The above tests are not answered by ketones.

## CARBOXYLIC ACIDS

Carbon compounds containing a carboxyl functional group (–COOH) are called carboxylic acids. The carboxyl group consists of a carbonyl group attached to a hydroxyl group, hence its name carboxyl. Aliphatic carboxylic acids containing 12 to 18 C atoms are called fatty acids. They occur in natural fats as esters of glycerol.

#### Nomenclature

The common names of carboxylic acids end with the suffix –ic acid and have been derived from Latin or Greek names of their natural sources. For example, formic acid (HCOOH) was first obtained from red ants (Latin: ormica means ant), acetic acid (CH<sub>3</sub>COOH) from vinegar (Latin: acetum, means vinegar), butyric acid (CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>COOH) from rancid butter (Latin: butyrum means butter).

In the IUPAC system, aliphatic carboxylic acids are named by replacing the ending –e in the name of the corresponding alkane with – oic acid.

In numbering the carbon chain, the carboxylic carbon is numbered one. For naming compounds containing more than one carboxyl group, the ending –e of the alkane is retained. The number of carboxyl groups is indicated by adding the prefixes di, tri, etc. to the term oic.

Some examples are:

Compound	Common name	IUPAC name
НСООН	Formic acid	Methanoic acid
CH <sub>3</sub> COOH	Acetic acid	Ethanoic acid
CH <sub>3</sub> CH <sub>2</sub> COOH	Propionic acid	Propanoic acid
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> COOH	Butyric acid	Butanoic acid
(CH <sub>3</sub> ) <sub>2</sub> CHCOOH	Isobutyric acid	2-Methylpropanoic acid
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> COOH	Valeric acid	Pentanoic acid
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> COOH	Caproic acid	Hexanoic acid
НООС-СООН	Oxalic acid	Ethanedioic acid
HOOC -CH <sub>2</sub> -COOH	Malonic acid	Propanedioic acid
HOOC -(CH <sub>2</sub> ) <sub>2</sub> -COOH	Succinic acid	Butanedioic acid
HOOC -(CH <sub>2</sub> ) <sub>3</sub> -COOH	Glutaric acid	Pentanedioic acid
HOOC -(CH <sub>2</sub> ) <sub>4</sub> -COOH	Adipic acid	Hexanedioic acid
HOOC -CH <sub>2</sub> -CH(COOH)-		Propane-1, 2, 3-
CH <sub>2</sub> -COOH		tricarboxylic acid
	Benzoic acid	Benzenecarboxylic acid
Соон		(Benzoic acid)
CH <sub>2</sub> COOH	Phenylacetic acid	2-Phenylethanoic acid
COOH		Benzene-1, 2-
Соон	Phthalic acid	dicarboxylic acid