

CHEMISTRY STUDY MATERIALS FOR CLASS 12

(NCERT Based Notes of Chapter - 13)

GANESH KUMAR

DATE:- 22/11/2020

Amines

Chemical Reactions

5. Reaction with nitrous acid:

Primary aliphatic amines react with nitrous acid (prepared by mixing sodium nitrite and HCl) to form alcohols with the liberation of nitrogen gas. From the amount of nitrogen evolved, we can estimate amino acids and proteins.



Primary amine

Diazonium salt

Alcohol

Aromatic primary amines react with nitrous acid at 0 to 5°C (273-278 K) to form aromatic diazonium salts.




Aniline

Benzene Diazonium chloride

6. Reaction with benzene sulphonyl chloride [Hinsberg's Test]:

This test is used for the distinction of primary, secondary and tertiary amines and also for the separation of a mixture of amines. Benzene sulphonyl chloride (C₆H₅SO₂Cl) is known as Hinsberg's reagent.

a) Primary amines react with benzene sulphonyl chloride to form N-alkyl benzene sulphonamide, which is soluble in alkali.




Benzene
sulphonyl
chloride

Ethanamine

N-Ethyl benzene sulphonamide

- b) Secondary amines react with benzene sulphonyl chloride to give N, N- dialkyl benzene sulphonamide, which is insoluble in alkali.



Benzene sulphonyl chloride + N- ethyl ethanamine → N,N-diethyl benzene sulphonamide + HCl

- c) Tertiary amines do not react with benzenesulphonyl chloride.

Nowadays, Benzene sulphonyl chloride is replaced by p-toluene sulphonyl chloride.

7. Electrophilic substitution Reactions:


–NH₂ group is ortho and para directing and a powerful activating group.

So the reactions are very vigorous.

- a) **Bromination:** Aniline reacts with bromine water at room temperature to give a white precipitate of 2,4,6-tribromoaniline.




In order to prepare o-bromoaniline and p-bromoaniline, first reduce the activating power of –NH₂ group by protecting it through acetylation with acetyl chloride or acetic anhydride. Then the resulting acetanilide is brominated by Br₂ in acetic acid followed by hydrolysis, we get p-bromoaniline as the major product.




b) Nitration:

Direct nitration of aniline with conc. HNO_3 and conc. H_2SO_4 gives a mixture of ortho, meta and para nitro anilines and some tarry (tar-like) products.




In strongly acidic medium, aniline is protonated to form the anilinium ion which is meta directing. So a large amount of meta-isomer is formed.

For the preparation of p-nitro aniline, the $-\text{NH}_2$ group is first deactivated by acetylation. The acetanilide thus formed is nitrated followed by hydrolysed.



c) Sulphonation:

Aniline reacts with concentrated sulphuric acid to form anilinium hydrogensulphate which on heating with sulphuric acid at 453-473K produces *p*-amino benzene sulphonic acid, commonly known as sulphanilic acid, as the major product. Sulphanilic acid contains both acidic and basic groups and so it forms internal salts called zwitter ions.



Aniline does not undergo Friedel-Crafts reaction (alkylation and acylation) since it forms salt with anhydrous aluminium chloride, which is used as catalyst in the reaction.
